

GEOMETRIC AUSLANDER CRITERION FOR FLATNESS OF AN ANALYTIC MAPPING

JANUSZ ADAMUS, EDWARD BIERSTONE AND PIERRE D. MILMAN

ABSTRACT. We prove that, if $\mathcal{F}$ is a coherent sheaf of $\mathcal{O}_X$ -modules over a morphism $\varphi : X \rightarrow Y$ of complex-analytic spaces, where Y is smooth, then the stalk $\mathcal{F}_\xi$ at a point $\xi \in X$ is flat over $R := \mathcal{O}_{Y, \varphi(\xi)}$ if and only if the n -fold analytic tensor power of $\mathcal{F}_\xi$ over R (where $n = \dim R$) has no *vertical elements*. The result implies that if $\mathcal{F}$ is a finite module over a morphism $\varphi : X \rightarrow Y$ of complex algebraic varieties, where Y is smooth and $\dim Y = n$, then $\mathcal{F}_\xi$ is R -flat if and only if its n -fold tensor power is a torsionfree R -module. The latter generalizes a classical freeness criterion of Auslander to modules that are not necessarily finitely generated over the base ring.

CONTENTS

1. Introduction	1
2. Analytic tensor product and fibred product	7
3. Homological properties of almost finitely generated modules	9
4. Vertical components and variation of fibre dimension	12
5. Proof of the main theorem	13
References	20

1. INTRODUCTION

This work is of origin in M. Auslander's criterion for freeness of a finitely generated module over a regular local ring [3, Thm. 3.2]:

Key words and phrases. flat, torsion free, fibred power, vertical component, analytic tensor product, complex analytic geometry.

Research partially supported by Natural Sciences and Engineering Research Council of Canada Discovery Grant OGP 355418-2008 and Polish Ministry of Science Discovery Grant N201 020 31/1768 (Adamus), and NSERC Discovery Grants OGP 0009070 (Bierstone), OGP 0008949 (Milman).

Theorem 1.1. *Let R be a regular local ring of dimension $n > 0$, and let F be a finite R -module. Then F is R -free if and only if the n -fold tensor power $F^{\otimes_R^n}$ is a torsion-free R -module.*

(Theorem 1.1 was proved in the case that R is unramified by Auslander, and extended to arbitrary regular local rings by Lichtenbaum [14].)

Our main result is an effective geometric analogue of Auslander's criterion, for flatness of a coherent sheaf $\mathcal{F}$ of $\mathcal{O}_X$ -modules over a morphism $\varphi : X \rightarrow Y$ of complex-analytic spaces, where Y is smooth (Theorem 1.10 below.) The result is an extension of a theorem of Galligo and Kwieciński [8].

In the context of Auslander's theorem, flat is equivalent to free. Vasconcelos [17] and Kwieciński [13] were the first to consider extending Auslander's criterion from finite modules to finite algebras over a regular local ring. Vasconcelos made the following conjecture [17, Conj. 6.2], [18, Conj. 2.6.1].

Conjecture 1.2. *Let R be a regular ring of dimension $n > 0$ and let A denote an R -algebra essentially of finite type. Then A is R -flat if and only if $A^{\otimes_R^n}$ is torsion-free over R .*

An R -algebra *essentially of finite type* means a localization of an R -algebra of finite type. Vasconcelos proved the analogue of Conjecture 1.2 for a Noetherian R -algebra over a 2-dimensional regular local ring R [17, Prop. 6.1].

We use our main theorem to prove a generalization of Vasconcelos's conjecture in the case that R is a $\mathbb{C}$ -algebra (see §1.2):

Theorem 1.3. *Let R be a regular $\mathbb{C}$ -algebra of finite type. Let A denote an R -algebra of finite type and let F denote a finitely generated A -module. Set $n = \dim R$. If $\mathfrak{p} \subset A$ is a prime ideal, then the localization $F_{\mathfrak{p}}$ is R -flat if and only if the n -fold tensor power $F_{\mathfrak{p}}^{\otimes_R^n}$ is a torsion-free R -module.*

Equivalently, if $\mathfrak{p} \subset A$ is a prime ideal and $\mathfrak{q} = \mathfrak{p} \cap R$, then $F_{\mathfrak{p}}$ is $R_{\mathfrak{q}}$ -flat if and only if $F_{\mathfrak{p}}^{\otimes_R^n}$ is $R_{\mathfrak{q}}$ -torsion-free.

Corollary 1.4. *With the assumptions of Theorem 1.3, F is R -flat if and only if $F^{\otimes_R^n}$ is R -torsion-free.*

We work with $\mathbb{C}$ -algebras (or, more generally, algebras over a complete algebraically-closed field) in this article because an important transcendental technique — Frisch's generic flatness theorem [7, Prop. VI,14] — plays an essential part (in the proof of *verticality* of torsion modules [8, Prop. 4.5]; see Lemma 3.1(4) below).

We are planning to write a sequel to this article, *Auslander criterion for flatness of a morphism of algebraic varieties*, in which we will use Grothendieck's generic flatness theorem [10, Thm. 11.1.1] rather than Frisch's theorem to prove the assertion of Theorem 1.3 in the case that R is a regular $\underline{k}$ -algebra of finite type over a field $\underline{k}$. This result establishes a generalization of Vasconcelos's conjecture in a natural algebraic-geometry setting — it provides a criterion for flatness of a coherent sheaf $\mathcal{F}$ of $\mathcal{O}_X$ -modules over a morphism $\varphi : X \rightarrow Y$ of schemes of finite type over $\underline{k}$.

In a second paper to follow, *An inductive geometric criterion for flatness*, we will prove another geometric flatness criterion using techniques which differ from but share a common approach with those here — successive reduction in fibre dimension via Weierstrass preparation, to eventually reduce the problem to a question about flatness of finitely-generated modules that is well-understood. The inductive criterion is a convenient tool to prove generic flatness results like Frisch's theorem.

Remark 1.5. Let R be a ring and let F be a finitely-generated module over an R -algebra A of finite type. From a computational point of view, a general way to investigate R -flatness of F is to find an element $r \in R$, such that (i) r is a zero-divisor in neither R nor F , and (ii) the localization R_r is regular (see [18]). Condition (i) permits reduction of dimension by passage from R to $R/(r)$. To exploit (ii), we need to handle the case of a regular ring directly; for example, using Theorem 1.3. The latter together with algorithms for primary decomposition (see [18]) provides an effective criterion for flatness in the case that R is a regular $\mathbb{C}$ -algebra of finite type: By Theorem 1.3 and the prime avoidance lemma [4, Lemma 3.3], in order to verify that F is not R -flat, it is enough to find an associated prime of $F^{\otimes_R^n}$ in $A^{\otimes_R^n}$ which contains a nonzero element $r \in R$. Note that n is independent of F . The power $n = \dim R$ is best possible already in the case that F is a finite R -module [3, §3], [18, Remark 2.62].

1.1. Vertical components in fibred powers. Kwieciński [13] proved that, if R is a finitely generated $\mathbb{C}$ -algebra which is a normal domain and A is a finitely generated R -algebra, then A is R -flat if and only if *all* tensor powers $A^{\otimes_R^k}$ are R -torsion-free. He used techniques of complex-analytic geometry, introducing the idea of a *vertical component* of an analytic mapping as a geometric analogue of torsion in commutative algebra. Following [1], we distinguish algebraic and geometric versions of Kwieciński's idea: Let $\varphi_\xi : X_\xi \rightarrow Y_\eta$ denote a morphism of germs of complex-analytic spaces.

Definition 1.6. Let W_ξ denote an irreducible component of X_ξ (isolated or embedded). We say that W_ξ is an *algebraic* (respectively, *geometric*) *vertical component* of φ_ξ (or over Y_η) if φ_ξ maps W_ξ to a proper analytic (respectively, nowhere-dense) subgerm of Y_η .

(We are allowing ourselves some imprecision of language. The notation $\varphi_\xi : X_\xi \rightarrow Y_\eta$ is meant to suggest the germ at a point $\xi \in X$ of a morphism of complex-analytic spaces $\varphi : X \rightarrow Y$, $\eta = \varphi(\xi)$. In particular, we will write $\mathcal{O}_{X,\xi}$ for the local ring of X_ξ . Consider a representative $\varphi : X \rightarrow Y$ of φ_ξ . The “if” clause in Definition 1.6 means more precisely that, for a sufficiently small representative W of W_ξ in X , the germ $\varphi(W)_\eta$ lies in a proper analytic subgerm of Y_η (respectively, $\varphi(W)$ is nowhere-dense in Y .)

By the prime avoidance lemma, $\varphi_\xi : X_\xi \rightarrow Y_\eta$ has an algebraic (respectively, geometric) vertical component if and only if there exists a nonzero element $m \in \mathcal{O}_{X,\xi}$ such that the zero-set germ $\mathcal{V}(\text{Ann}_{\mathcal{O}_{X,\xi}}(m))$ of the annihilator of m in $\mathcal{O}_{X,\xi}$ is mapped to a proper analytic (respectively, nowhere-dense) subgerm of Y_η .

We can extend the notion of vertical component to a finitely generated $\mathcal{O}_{X,\xi}$ -module F :

Definition 1.7. Let Z_ξ be the zero-set germ of the support of F ; i.e., $Z_\xi = \mathcal{V}(\bigcap\{\mathfrak{p} : \mathfrak{p} \in \text{Ass}_{\mathcal{O}_{X,\xi}}(F)\})$. We say that F has an *algebraic* (respectively, *geometric*) *vertical component* over Y_η (or over $\mathcal{O}_{Y,\eta}$) if Z_ξ has an algebraic (respectively, geometric) vertical component over Y_η in the sense of Definition 1.6; equivalently (by prime avoidance again), there exists a nonzero $m \in F$ such that the $\mathcal{V}(\text{Ann}_{\mathcal{O}_{X,\xi}}(m))$ is mapped to a proper analytic (respectively, nowhere-dense) subgerm of Y_η . In the geometric case, we will call such m a *geometric vertical element* (or simply a *vertical element*) of F over Y_η (or over $\mathcal{O}_{Y,\eta}$).

Note that an analogous “algebraic vertical element” of F over Y_η is simply a (nonzero) zero-divisor of F over $\mathcal{O}_{Y,\eta}$, so there is no need to define algebraic vertical elements. A *vertical element* will always mean geometric vertical.

Remark 1.8. In the special case that $F = \mathcal{O}_{X,\xi}$, X_ξ has no geometric (respectively, algebraic) vertical components over Y_η if and only if $\mathcal{O}_{X,\xi}$ (as an $\mathcal{O}_{X,\xi}$ -module) has no vertical elements (respectively, no zero-divisors) over $\mathcal{O}_{Y,\eta}$.

Now let R denote a regular local analytic $\mathbb{C}$ -algebra of dimension n . Then R is isomorphic to the ring $\mathbb{C}\{y\} = \mathbb{C}\{y_1, \dots, y_n\}$ of convergent power series in n variables. A *local analytic R -algebra* A

means a ring of the form $R\{x\}/I = \mathbb{C}\{y, x\}/I$, where I is an ideal in $\mathbb{C}\{y, x\} = \mathbb{C}\{y_1, \dots, y_n, x_1, \dots, x_m\}$, with the canonical homomorphism $R \rightarrow A$. Let F denote an R -module. We say that F is an *almost finitely generated R -module* (following [8]) if F is a finitely-generated A -module, for some local analytic R -algebra A . In this case, there is a morphism of germs of analytic spaces $\varphi_\xi : X_\xi \rightarrow Y_\eta$ such that $R \cong \mathcal{O}_{Y,\eta}$, $A \cong \mathcal{O}_{X,\xi}$, $R \rightarrow A$ is the induced homomorphism $\varphi_\xi^* : \mathcal{O}_{Y,\eta} \rightarrow \mathcal{O}_{X,\xi}$, and F is a finitely-generated $\mathcal{O}_{X,\xi}$ -module. We say that a nonzero element $m \in F$ is *vertical* over R if m is vertical over $\mathcal{O}_{Y,\eta}$ in the sense of Definition 1.7.

Remark 1.9. It is easy to see that the notion of vertical element is well-defined; i.e., independent of a choice of local R -algebra A such that F is a finitely generated A -module. In particular, given an almost finitely generated R -module F , we can assume without loss of generality that F is finitely generated over the regular ring $A = R\{x\} \cong \mathbb{C}\{y, x\}$, where $x = (x_1, \dots, x_m)$, for some $m \geq 0$.

1.2. Main results. Our main theorem is the following flatness criterion.

Theorem 1.10. *Let R be a regular local analytic $\mathbb{C}$ -algebra and let F denote an almost finitely generated R -module. Let $n = \dim R$. Then F is R -flat if and only if the n -fold analytic tensor power $F^{\tilde{\otimes}_R^n}$ has no vertical elements over R .*

(See Section 2 for the notion of analytic tensor power.) In the case that F is a local analytic R -algebra A , Theorem 1.10 can be restated as follows.

Corollary 1.11. *Let $\varphi_\xi : X_\xi \rightarrow Y_\eta$ denote a morphism of germs of complex-analytic spaces, where Y_η is smooth of dimension n . Then φ_ξ is flat if and only if the n -fold fibred power $\varphi_{\xi\{n\}}^{\{n\}} : X_{\xi\{n\}}^{\{n\}} \rightarrow Y_\eta$ has no (isolated or embedded) geometric vertical components.*

Proof. Let $R = \mathbb{C}\{y_1, \dots, y_n\}$. Then $\mathcal{O}_{Y,\eta} \cong R$ and $\mathcal{O}_{X,\xi}$ can be viewed as an almost finitely generated R -module. (We can embed (a representative of) X_ξ in $\mathbb{C}^m$ and identify X_ξ with the graph of φ_ξ in $\mathbb{C}_\xi^m \times Y_\eta$ to realize $\mathcal{O}_{X,\xi}$ as a quotient of $R\{x_1, \dots, x_m\}$.) Since $\mathcal{O}_{X,\xi} \tilde{\otimes}_{\mathcal{O}_{Y,\eta}} \dots \tilde{\otimes}_{\mathcal{O}_{Y,\eta}} \mathcal{O}_{X,\xi} \cong \mathcal{O}_{X\{n\},\xi\{n\}}$, the left-hand side has no vertical elements over $\mathcal{O}_{Y,\eta}$ if and only if the n -fold fibred power of X_ξ over Y_η has no geometric vertical components (by Remark 1.8). $\square$

Corollary 1.11 in the special case that X_ξ is equidimensional is the theorem of Galligo and Kwieciński [8]. The assumption that X_ξ is

equidimensional guarantees that $A = \mathcal{O}_{X,\xi}$ is a finite *torsion-free* module over some R -flat algebra S (where $R = \mathcal{O}_{Y,\eta}$), and Auslander's techniques can be extended to this case [8].

We are happy to acknowledge the influence of [8] on our paper. We show that the same methods can be used when the module F in Theorem 1.10 is torsion-free over A . (See Section 3 and Case (1) of the proof of Theorem 1.10.) Of course, if F is expressed as in Remark 1.9 and $m = 0$, Auslander's theorem applies directly. We prove Theorem 1.10 by induction of the number m of x -variables. The most difficult new situation in the inductive step is the case that F is neither torsion-free nor a torsion A -module (Case (3) of the proof of Theorem 1.10.)

By way of comparison with Corollary 1.11, we note that *openness* of an analytic mapping with irreducible target can be characterized by the lack of *isolated algebraic* vertical components in the n -fold fibred power (where n is the target dimension; see [1], [2], and Section 4).

Frisch's generic flatness theorem asserts that if $\mathcal{F}$ is a coherent sheaf of $\mathcal{O}_X$ -modules over a morphism $\varphi : X \rightarrow Y$ of complex-analytic spaces, then $\{\xi \in X : \mathcal{F}_\xi \text{ is not } \mathcal{O}_{Y,\varphi(\xi)}\text{-flat}\}$ has image nowhere-dense in Y . Frisch's theorem is responsible for the criterion of Theorem 1.10 in terms of *geometric* vertical elements. The existence of a vertical element in $F^{\otimes_R^n}$ guarantees the existence of a zero-divisor of $F^{\otimes_R^k}$ over R , for *some* $k \geq n$ (by [1] and Theorem 1.10). The following question seems to be open.

Question 1.12. *With the assumptions of Theorem 1.10, if F is not R -flat, does $F^{\otimes_R^n}$ have a zero-divisor over R ?*

If $\varphi_\xi : X_\xi \rightarrow Y_\eta$ and F have an underlying algebraic structure as in Theorem 1.3, then the notions of geometric and algebraic vertical components coincide, so that Theorem 1.3 follows from Theorem 1.10:

Proof of Theorem 1.3. If $F_{\mathfrak{p}}$ is R -flat, then $F_{\mathfrak{p}}^{\otimes_R^k}$ is R -flat and therefore R -torsion-free, for all k .

On the other hand, flatness and torsion-freeness are both local properties. I.e., $F_{\mathfrak{p}}$ is R -flat (respectively, R -torsion-free) if and only if $F_{\mathfrak{m}}$ is R -flat (respectively, R -torsion-free), for every maximal ideal $\mathfrak{m}$ of A containing $\mathfrak{p}$. Suppose that $F_{\mathfrak{p}}$ is not R -flat. Then there is a maximal ideal $\mathfrak{m}$ containing $\mathfrak{p}$ such that $F_{\mathfrak{m}}$ is not R -flat, and it suffices to prove that $F_{\mathfrak{m}}^{\otimes_R^n}$ is not R -torsion-free. Let $\varphi : X \rightarrow Y$ be the morphism of complex-analytic spaces associated to the morphism $\text{Spec } A \rightarrow \text{Spec } R$ and let $\mathcal{F}$ be the coherent sheaf of $\mathcal{O}_X$ -modules associated to F . Let $\xi \in X$ be the point corresponding to the maximal ideal $\mathfrak{m}$ of $\text{Spec } A$. It follows from faithful flatness of completion that $\mathcal{F}_\xi$ is not $\mathcal{O}_{Y,\eta}$ -flat,

where $\eta = \varphi(\xi)$. By Theorem 1.10, $\mathcal{F}_\xi^{\tilde{\otimes}_{\mathcal{O}_{Y,\eta}}^n}$ has a vertical element over $\mathcal{O}_{Y,\eta}$. Since $\varphi^{\{n\}}$ is the holomorphic map induced by the ring homomorphism $R \rightarrow A^{\otimes_R^n}$, it follows from Chevalley's Theorem that $\mathcal{F}_\xi^{\tilde{\otimes}_{\mathcal{O}_{Y,\eta}}^n}$ has a zero-divisor in $\mathcal{O}_{Y,\eta}$. Then $F_{\mathfrak{m}}^{\otimes_R^n}$ has a zero-divisor in R ; i.e., $F_{\mathfrak{m}}^{\otimes_R^n}$ is not R -torsion-free. $\square$

2. ANALYTIC TENSOR PRODUCT AND FIBRED PRODUCT

We briefly recall the concepts of analytic tensor product and fibred product of analytic spaces, which are used throughout the paper.

The analytic tensor product is defined in the category of finitely generated modules over local analytic $\mathbb{C}$ -algebras (i.e., rings of the form $\mathbb{C}\{z_1, \dots, z_n\}/I$ for some ideal I) by the usual universal mapping property for tensor product (cf. [9]): Let $\varphi_i : R \rightarrow A_i$ ($i = 1, 2$) be homomorphisms of local analytic $\mathbb{C}$ -algebras. Then there is a unique (up to isomorphism) local analytic $\mathbb{C}$ -algebra $A_1 \tilde{\otimes}_R A_2$, together with homomorphisms $\theta_i : A_i \rightarrow A_1 \tilde{\otimes}_R A_2$ ($i = 1, 2$), such that (1) $\theta_1 \circ \varphi_1 = \theta_2 \circ \varphi_2$, and (2) for every pair of homomorphisms of local analytic $\mathbb{C}$ -algebras $\psi_1 : A_1 \rightarrow B$, $\psi_2 : A_2 \rightarrow B$ satisfying $\psi_1 \circ \varphi_1 = \psi_2 \circ \varphi_2$, there is a unique homomorphism of local analytic $\mathbb{C}$ -algebras $\psi : A_1 \tilde{\otimes}_R A_2 \rightarrow B$ making the associated diagram commute. The algebra $A_1 \tilde{\otimes}_R A_2$ is called the *analytic tensor product* of A_1 and A_2 over R .

For finite modules M_1 and M_2 over local analytic R -algebras A_1 and A_2 , respectively, there is a unique (up to isomorphism) finite $A_1 \tilde{\otimes}_R A_2$ -module $M_1 \tilde{\otimes}_R M_2$, together with an R -bilinear mapping $\rho : M_1 \times M_2 \rightarrow M_1 \tilde{\otimes}_R M_2$, such that for every R -bilinear $\kappa : M_1 \times M_2 \rightarrow N$, where N is a finite $A_1 \tilde{\otimes}_R A_2$ -module, there is a unique homomorphism of $A_1 \tilde{\otimes}_R A_2$ -modules $\lambda : M_1 \tilde{\otimes}_R M_2 \rightarrow N$ satisfying $\kappa = \lambda \circ \rho$. The module $M_1 \tilde{\otimes}_R M_2$ is called the *analytic tensor product* of M_1 and M_2 over R .

Remark 2.1. The analytic tensor product is often denoted $\hat{\otimes}$, although there seems to be no standard notation. We use the notation $\tilde{\otimes}$ for the analytic tensor product (and $\widetilde{\text{Tor}}$ for its derived functors) in order to avoid confusion with the completed tensor product $\hat{\otimes}$ (and its derived functors) which also appear frequently in our arguments.

It is sometimes convenient to express the analytic tensor product of modules over a local analytic $\mathbb{C}$ -algebra in terms of ordinary tensor product of certain naturally associated modules: Given homomorphisms of local analytic $\mathbb{C}$ -algebras $\varphi : R \rightarrow A_i$, and finitely generated A_i -modules M_i ($i = 1, 2$), the modules $M_1 \tilde{\otimes}_R A_2$ and $A_1 \tilde{\otimes}_R M_2$ are

finitely generated over $A_1 \tilde{\otimes}_R A_2$, and there is a canonical isomorphism

$$M_1 \tilde{\otimes}_R M_2 \cong (M_1 \tilde{\otimes}_R A_2) \otimes_{A_1 \tilde{\otimes}_R A_2} (A_1 \tilde{\otimes}_R M_2).$$

In particular, if $A_1 = R\{x\}/I_1$ and $A_2 = R\{t\}/I_2$, where $x = (x_1, \dots, x_l)$, $t = (t_1, \dots, t_m)$ are systems of variables and $I_1 \subset R\{x\}$, $I_2 \subset R\{t\}$ are ideals, then

$$\begin{aligned} A_1 \tilde{\otimes}_R A_2 &\cong (A_1 \tilde{\otimes}_R R\{t\}) \otimes_{R\{x\} \tilde{\otimes}_R R\{t\}} (R\{x\} \tilde{\otimes}_R A_2) \\ &\cong (R\{x, t\}/I_1 R\{x, t\}) \otimes_{R\{x, t\}} (R\{x, t\}/I_2 R\{x, t\}) \\ &\cong R\{x, t\}/(I_1 R\{x, t\} + I_2 R\{x, t\}). \end{aligned}$$

The fibred product of analytic spaces is defined by a dual universal mapping property (see [6]): Let $\varphi : X_i \rightarrow Y$ ($i = 1, 2$) denote holomorphic mappings of complex analytic spaces. Then there exists a unique (up to isomorphism) complex analytic space $X_1 \times_Y X_2$, together with holomorphic maps $\pi_i : X_1 \times_Y X_2 \rightarrow X_i$ ($i = 1, 2$), such that (1) $\varphi_1 \circ \pi_1 = \varphi_2 \circ \pi_2$, and (2) for every pair of holomorphic maps $\psi_1 : X \rightarrow X_1$, $\psi_2 : X \rightarrow X_2$ satisfying $\varphi_1 \circ \psi_1 = \varphi_2 \circ \psi_2$, there is a unique holomorphic map $\psi : X \rightarrow X_1 \times_Y X_2$ making the associated diagram commute. The space $X_1 \times_Y X_2$ is called the *fibred product* of X_1 and X_2 over Y (more precisely, over φ_1 and φ_2). There is a canonical holomorphic mapping $\varphi_1 \times_Y \varphi_2 : X_1 \times_Y X_2 \rightarrow Y$, given by $\varphi_1 \times_Y \varphi_2 = \varphi_i \circ \pi_i$ (where $i = 1$ or 2).

Given a holomorphic map $\varphi : X \rightarrow Y$ of complex analytic spaces, with $\varphi(\xi) = \eta$, let $\varphi_\xi : X_\xi \rightarrow Y_\eta$ denote the germ of φ at ξ . We denote by $\varphi^{\{d\}} : X^{\{d\}} \rightarrow Y$ the canonical map from the d -fold fibred power of X over Y to Y , and by $\varphi_{\xi^{\{d\}}}^{\{d\}} : X_{\xi^{\{d\}}}^{\{d\}} \rightarrow Y_\eta$ its germ at the point $\xi^{\{d\}} := (\xi, \dots, \xi) \in X^d$.

Suppose that $\varphi_1 : X_1 \rightarrow Y$ and $\varphi_2 : X_2 \rightarrow Y$ are holomorphic mappings of analytic spaces, with $\varphi_1(\xi_1) = \varphi_2(\xi_2) = \eta$. Then the local rings $\mathcal{O}_{X_i, \xi_i}$ ($i = 1, 2$) are $\mathcal{O}_{Y, \eta}$ -modules and, by the uniqueness of fibred product and of analytic tensor product, the local ring $\mathcal{O}_{Z, (\xi_1, \xi_2)}$ of the fibred product $Z = X_1 \times_Y X_2$ at (ξ_1, ξ_2) is canonically isomorphic to $\mathcal{O}_{X_1, \xi_1} \tilde{\otimes}_{\mathcal{O}_{Y, \eta}} \mathcal{O}_{X_2, \xi_2}$. Therefore, given a holomorphic germ $\varphi_\xi : X_\xi \rightarrow Y_\eta$, we will identify the d -fold analytic tensor power $\mathcal{O}_{X, \xi}^{\tilde{\otimes}^d \mathcal{O}_{Y, \eta}} = \mathcal{O}_{X, \xi} \tilde{\otimes}_{\mathcal{O}_{Y, \eta}} \dots \tilde{\otimes}_{\mathcal{O}_{Y, \eta}} \mathcal{O}_{X, \xi}$ with the local ring of the d -fold fibred power $\mathcal{O}_{X^{\{d\}}, \xi^{\{d\}}}$, for $d \geq 1$.

3. HOMOLOGICAL PROPERTIES OF ALMOST FINITELY GENERATED MODULES

We begin with some flatness properties of almost finitely generated modules, established by Galligo and Kwieciński [8], that generalize the corresponding properties of finite modules used by Auslander's [3]. We then generalize a lemma of Auslander [3, Lemma 3.1] to almost finitely generated modules (Lemma 3.3 below).

Let $R = \mathbb{C}\{y_1, \dots, y_n\}$ denote a regular local analytic $\mathbb{C}$ -algebra of dimension n . Let $\tilde{\otimes}_R$ denote the analytic tensor product over R , and let $\widetilde{\mathrm{Tor}}^R$ be the corresponding derived functor.

Let F denote an almost finitely generated R -module. We define the *flat dimension* $\mathrm{fd}_R(F)$ of F over R as the minimal length of a flat resolution of F (i.e., a resolution by R -flat modules). It is easy to see that

$$(3.1) \quad \mathrm{fd}_R(F) = \max\{i \in \mathbb{N} : \widetilde{\mathrm{Tor}}_i^R(F, N) \neq 0 \text{ for some } N\}.$$

Indeed, if M is an almost finitely generated R -module, then

$$(3.2) \quad M \text{ is } R\text{-flat} \Leftrightarrow \widetilde{\mathrm{Tor}}_1^R(M, R/\mathfrak{m}_R) = 0,$$

where $\mathfrak{m}_R$ is the maximal ideal of R (cf. [11, Prop. 6.2]). Let $(A, \mathfrak{m}_A)$ be a regular local R -algebra such that F is a finite A -module. Then (3.1) follows from (3.2) applied to the kernels of a minimal A -free (hence R -flat) resolution

$$\mathcal{F}_* : \quad \dots \xrightarrow{\alpha_{i+1}} F_{i+1} \xrightarrow{\alpha_i} F_i \xrightarrow{\alpha_{i-1}} \dots \xrightarrow{\alpha_1} F_1 \xrightarrow{\alpha_0} F_0 \rightarrow F$$

of F . ($\mathcal{F}_*$ *minimal* means that $\alpha_i(F_{i+1}) \subset \mathfrak{m}_A F_i$, for all $i \in \mathbb{N}$).

The *depth* $\mathrm{depth}_R(F)$ of F as an R -module is defined as the length of a maximal F -sequence in R (i.e., a sequence $a_1, \dots, a_s \in \mathfrak{m}_R$ such that a_j is not a zero-divisor in $F/(a_1, \dots, a_{j-1})F$, for $j = 1, \dots, s$). Since all the maximal F -sequences in R have the same length, depth is well defined: As observed in [8, Lemma 2.4], the classical proof of Northcott-Rees for finitely generated modules (see, e.g., [12, §VI, Prop. 3.1]), carries over to the case of almost finitely generated modules.

Lemma 3.1. *Let M and N be almost finitely generated R -modules. Then the following properties hold.*

- (1) Rigidity of $\widetilde{\mathrm{Tor}}^R$ [8, Prop. 2.2(4)]. If $\widetilde{\mathrm{Tor}}_{i_0}^R(M, N) = 0$ for some $i_0 \in \mathbb{N}$, then $\widetilde{\mathrm{Tor}}_i^R(M, N) = 0$ for all $i \geq i_0$.
- (2) Auslander–Buchsbaum-type formula [8, Thm. 2.7].

$$\mathrm{fd}_R(M) + \mathrm{depth}_R(M) = n.$$

- (3) Additivity of flat dimension [8, Prop. 2.10]. If $\widetilde{\mathrm{Tor}}_i^R(M, N) = 0$ for all $i \geq 1$, then

$$\mathrm{fd}_R(M) + \mathrm{fd}_R(N) = \mathrm{fd}_R(M \tilde{\otimes}_R N).$$

- (4) Verticality of $\widetilde{\mathrm{Tor}}^R$ (cf. [8, Prop. 4.5]). For all $i \geq 1$, $\widetilde{\mathrm{Tor}}_i^R(M, N)$ is an almost finitely generated R -module, and every element of $\widetilde{\mathrm{Tor}}_i^R(M, N)$ is vertical over R (recall Definition 1.7).

Remark 3.2. Contrary to the finitely generated case, the analytic $\widetilde{\mathrm{Tor}}_i^R$ need not be torsion R -modules. It seems to be unknown whether they necessarily contain R -zero-divisors (cf. Question 1.12).

Lemma 3.3. Let $A = R\{x\}$ denote a regular local analytic R -algebra, $x = (x_1, \dots, x_m)$. Let F be a finitely generated A -torsion-free module, and let N be a module which is finitely generated over $B = A^{\tilde{\otimes}_R^j}$, for some $j \geq 1$. Suppose that $F \tilde{\otimes}_R N$ has no vertical elements over R . Then:

- (1) N has no vertical elements over R ;
- (2) $\widetilde{\mathrm{Tor}}_i^R(F, N) = 0$, for all $i \geq 1$;
- (3) $\mathrm{fd}_R(F) + \mathrm{fd}_R(N) = \mathrm{fd}_R(F \tilde{\otimes}_R N)$.

Proof. To prove (1), consider $N' = \{n \in N : n \text{ is vertical over } R\}$. It is easy to see that N' is a B -submodule of N . Indeed, if $n, n_1, n_2 \in N'$ and $b \in B$, then

$$\mathrm{Ann}_B(n_1 + n_2) \supset \mathrm{Ann}_B(n_1) \cdot \mathrm{Ann}_B(n_2) \quad \text{and} \quad \mathrm{Ann}_B(bn) \supset \mathrm{Ann}_B(n);$$

hence the zero set germ $\mathcal{V}(\mathrm{Ann}_B(n_1 + n_2))$ is mapped into the union of the (nowhere-dense) images of $\mathcal{V}(\mathrm{Ann}_B(n_1))$ and $\mathcal{V}(\mathrm{Ann}_B(n_2))$, and $\mathcal{V}(\mathrm{Ann}_B(bn))$ is mapped into the image of $\mathcal{V}(\mathrm{Ann}_B(n))$. Therefore, we get an exact sequence of B -modules,

$$0 \rightarrow N' \rightarrow N \rightarrow N'' = N/N' \rightarrow 0.$$

Tensoring with F induces a long exact sequence of $A^{\tilde{\otimes}_R^{j+1}}$ -modules,

$$\begin{aligned} (3.3) \quad \dots &\rightarrow \widetilde{\mathrm{Tor}}_{i+1}^R(F, N') \rightarrow \widetilde{\mathrm{Tor}}_{i+1}^R(F, N) \rightarrow \widetilde{\mathrm{Tor}}_{i+1}^R(F, N'') \\ &\rightarrow \widetilde{\mathrm{Tor}}_i^R(F, N') \rightarrow \dots \rightarrow \widetilde{\mathrm{Tor}}_1^R(F, N'') \\ &\rightarrow F \tilde{\otimes}_R N' \rightarrow F \tilde{\otimes}_R N \rightarrow F \tilde{\otimes}_R N'' \rightarrow 0. \end{aligned}$$

Since every element of N' is vertical over R , the same is true for $F \tilde{\otimes}_R N'$ (indeed, $\mathrm{Ann}_{A \tilde{\otimes}_R B}(f \tilde{\otimes}_R n) \supset 1 \tilde{\otimes}_R \mathrm{Ann}_B(n)$ for all $f \in F, n \in N'$). But $F \tilde{\otimes}_R N$ has no vertical elements, by assumption, so that $F \tilde{\otimes}_R N' \rightarrow$

$F \tilde{\otimes}_R N$ is the zero map; hence $F \tilde{\otimes}_R N \cong F \tilde{\otimes}_R N''$. In particular, $F \tilde{\otimes}_R N''$ has no vertical elements over R .

Since F is A -torsion-free, there is an injection of F into a finite free A -module L (obtained by composing the natural map $F \rightarrow (F^*)^*$, which is injective in this case, with the dual of a presentation of F^*). The exact sequence of A -modules $0 \rightarrow F \rightarrow L \rightarrow L/F \rightarrow 0$ induces a long exact sequence of $A \tilde{\otimes}_R B$ -modules,

$$\begin{aligned} \dots &\rightarrow \widetilde{\text{Tor}}_{i+1}^R(L, N'') \rightarrow \widetilde{\text{Tor}}_{i+1}^R(L/F, N'') \rightarrow \widetilde{\text{Tor}}_i^R(F, N'') \\ &\rightarrow \widetilde{\text{Tor}}_i^R(L, N'') \rightarrow \dots \rightarrow \widetilde{\text{Tor}}_1^R(L, N'') \rightarrow \widetilde{\text{Tor}}_1^R(L/F, N'') \\ &\rightarrow F \tilde{\otimes}_R N'' \rightarrow L \tilde{\otimes}_R N'' \rightarrow L/F \tilde{\otimes}_R N'' \rightarrow 0. \end{aligned}$$

Since L is a free A -module and therefore R -flat, $\widetilde{\text{Tor}}_i^R(L, N'') = 0$ for all $i \geq 1$, and we obtain isomorphisms

$$(3.4) \quad \widetilde{\text{Tor}}_{i+1}^R(L/F, N'') \cong \widetilde{\text{Tor}}_i^R(F, N''), \quad i \geq 1,$$

as well as injectivity of $\widetilde{\text{Tor}}_1^R(L/F, N'') \rightarrow F \tilde{\otimes}_R N''$. But $F \tilde{\otimes}_R N''$ has no vertical elements, while every element of $\widetilde{\text{Tor}}_1^R(L/F, N'')$ is vertical over R (by Lemma 3.1(4)); hence $\widetilde{\text{Tor}}_1^R(L/F, N'') \rightarrow F \tilde{\otimes}_R N''$ is the zero map. Therefore, $\dots \xrightarrow{0} \widetilde{\text{Tor}}_1^R(L/F, N'') \xrightarrow{0} \dots$ is exact; hence $\widetilde{\text{Tor}}_1^R(L/F, N'') = 0$. By rigidity of $\widetilde{\text{Tor}}^R$ (Lemma 3.1(1)), $\widetilde{\text{Tor}}_{i+1}^R(L/F, N'') = 0$ for all $i \geq 1$, so by (3.4),

$$(3.5) \quad \widetilde{\text{Tor}}_i^R(F, N'') = 0, \quad i \geq 1.$$

In particular, $\widetilde{\text{Tor}}_1^R(F, N'') = 0$, hence, $\dots \xrightarrow{0} F \tilde{\otimes}_R N' \xrightarrow{0} \dots$ is exact, by (3.3), so that $F \tilde{\otimes}_R N' = 0$. However, $F \tilde{\otimes}_R N' \cong (F \tilde{\otimes}_R B) \otimes_{A \tilde{\otimes}_R B} (A \tilde{\otimes}_R N')$ is an (ordinary) tensor product of finitely generated modules over a regular local ring $A \tilde{\otimes}_R B$, so it is zero only if one of the factors is zero. We conclude that $A \tilde{\otimes}_R N' = 0$, and therefore $N' = 0$, by R -flatness of A . This proves assertion (1).

Now, $\widetilde{\text{Tor}}_i^R(F, N') = 0$ for all $i \geq 0$; hence $\widetilde{\text{Tor}}_i^R(F, N) \cong \widetilde{\text{Tor}}_i^R(F, N'')$ for all $i \geq 1$, by (3.3). Therefore, $\widetilde{\text{Tor}}_i^R(F, N) = 0$ for all $i \geq 1$, by (3.5), proving (2).

Assertion (3) follows from Lemma 3.1(3) and (2). $\square$

4. VERTICAL COMPONENTS AND VARIATION OF FIBRE DIMENSION

In this section, we describe a relationship between the filtration of the target of an analytic mapping $\varphi : X \rightarrow Y$ by fibre dimension and the isolated irreducible components of the n -fold fibred power $X^{\{n\}}$, where $n = \dim Y$.

Let $\varphi_\xi : X_\xi \rightarrow Y_\eta$ be a morphism of germs of analytic spaces, where Y_η is irreducible and of dimension n . Let Y be an irreducible representative of Y_η , and let X be a representative of X_ξ , such that the components of X are precisely the representatives in X of the components of X_ξ , and $\varphi(X) \subset Y$, where φ represents the germ φ_ξ . Let $\text{fbd}_x \varphi$ denote the *fibre dimension* $\dim_x \varphi^{-1}(\varphi(x))$ of φ at a point $x \in X$.

We will use the following notation in this section: $l := \min\{\text{fbd}_x \varphi : x \in X\}$, $k := \max\{\text{fbd}_x \varphi : x \in X\}$, and $A_j := \{x \in X : \text{fbd}_x \varphi \geq j\}$, $l \leq j \leq k$. Then $X = A_l \supset A_{l+1} \supset \cdots \supset A_k$ and, by upper-semicontinuity of fibre dimension (see Cartan–Remmert Theorem [15, §V.3.3, Thm. 5]), the A_j are analytic in X . Define $B_j := f(A_j) = \{y \in Y : \dim \varphi^{-1}(y) \geq j\}$, $l \leq j \leq k$. Upper-semicontinuity of $\text{fbd}_x \varphi$ (as a function of x) implies that the germs $(A_j)_\xi$ and $(B_j)_\eta$ are independent of the choices of representatives made above.

Note that, except for B_k (cf. proof of Proposition 4.1 below), the B_j may not even be semianalytic in general. This fact is responsible for a complicated relationship between the algebraic vertical and geometric vertical components in the fibred powers of X over Y (see [2] for a detailed discussion), but will not affect our considerations here, which rely only on the properties of B_k .

Proposition 4.1 ([1, Prop.2.1]). *Under the assumptions above, let $\bigcup_{i \in I} W_i$ denote the decomposition of $(X^{\{n\}})_{\text{red}}$ into finitely many isolated irreducible components through $\xi^{\{n\}}$. Then:*

- (1) *For each $j = l, \dots, k$, there is an index subset $I_j \subset I$ such that*

$$B_j = \bigcup_{i \in I_j} \varphi^{\{n\}}(W_i).$$

- (2) *Let $y \in B_j$ and let $s = \dim \varphi^{-1}(y)$ ($s \geq j$). If Z is an isolated irreducible component of the fibre $(\varphi^{\{n\}})^{-1}(y)$, of dimension ns , and W is an irreducible component of $X^{\{n\}}$ containing Z , then $\varphi^{\{n\}}(W) \subset B_j$.*

Proof. For (2), fix $j \geq l + 1$. (The statement is trivial for $j = l$, since $B_l = \varphi(X)$.) Suppose that there exists $x = (x_1, \dots, x_n) \in W$ such that $\varphi(x_1) \in Y \setminus B_j$ (and hence $\varphi(x_i) \in Y \setminus B_j$, $i \leq n$). Then $\text{fbd}_{x_i} \varphi \leq j - 1$, $i = 1, \dots, n$; hence $\text{fbd}_x \varphi^{\{n\}} \leq n(j - 1) = nj - n$.

In particular, the generic fibre dimension of $\varphi^{\{n\}}|_W$ is at most $nj - n$. Since $\text{rank}(\varphi^{\{n\}}|_W) \leq \dim Y = n$, then $\dim W \leq (nj - n) + n = nj$ (see, e.g., [15, V.3]).

Now we have $W \supset Z$, $\dim W \leq nj$, $\dim Z = ns \geq nj$, and both W and Z are irreducible analytic sets in $X^{\{n\}}$. This is possible only if $W = Z$; hence $\varphi^{\{n\}}(W) = \varphi^{\{n\}}(Z) = \{y\} \subset B_j$; a contradiction. Therefore $\varphi^{\{n\}}(W) \subset B_j$, completing the proof of (2).

Part (1) follows immediately, since if $y \in B_j$ and Z is an irreducible component of $(\varphi^{\{n\}})^{-1}(y)$ of the highest dimension, then there exists an isolated irreducible component W of $X^{\{n\}}$ that contains Z . $\square$

The following is a simplified variant of an openness criterion of [1, Thm. 2.2], proved here under somewhat weaker assumptions.

Proposition 4.2. *Let $\varphi_\xi : X_\xi \rightarrow Y_\eta$ be a morphism of germs of analytic spaces. Suppose that Y_η is irreducible, $\dim Y_\eta = n$, $\dim X_\xi = m$, and the maximal fibre dimension of φ_ξ is not generic on any m -dimensional irreducible component of X_ξ . Then the n -fold fibred power $\varphi_{\xi^{\{n\}}}^{\{n\}} : X_{\xi^{\{n\}}}^{\{n\}} \rightarrow Y_\eta$ contains an isolated algebraic vertical component.*

Proof. As above, let $\varphi : X \rightarrow Y$ be a representative of φ_ξ , where Y is irreducible and of dimension n . Let $k := \max\{\text{fbd}_x \varphi : x \in X\}$, $A_k := \{x \in X : \text{fbd}_x \varphi = k\}$, and $B_k := \varphi(A_k) = \{y \in Y : \dim \varphi^{-1}(y) = k\}$. Then the fibre dimension of φ is constant on the analytic set A_k . By the Remmert Rank Theorem (see [15, §V.6, Thm. 1]), B_k is locally analytic in Y , of dimension $\dim X - k$. Shrinking Y if necessary, we can assume that B_k is an analytic subset of Y . Therefore, by Proposition 4.1, it is enough to show that the analytic germ $(B_k)_\eta$ is a proper subgerm of Y_η . Let U be an isolated irreducible component of X , of dimension $m = \dim X$. By assumption, the generic fibre dimension of $\varphi|_U$ is at most $k - 1$; hence

$$\dim Y \geq \dim U - \text{generic fbd} \varphi|_U \geq m - k + 1.$$

Then $\dim B_k = m - k < \dim Y$; hence $\dim(B_k)_\eta < \dim Y = \dim Y_\eta$, so that $(B_k)_\eta \subsetneq Y_\eta$. $\square$

5. PROOF OF THE MAIN THEOREM

Proof of Theorem 1.10. We can assume that F is an almost finitely generated module over $R := \mathbb{C}\{y_1, \dots, y_n\}$. By Remark 1.9, there exists $m \geq 0$ such that F is finitely generated as a module over $A = R\{x\}$, where $x = (x_1, \dots, x_m)$. Let X and Y be connected open neighbourhoods of the origins in $\mathbb{C}^{m+n}$ and $\mathbb{C}^n$ (respectively), and let $\varphi : X \rightarrow Y$

be the canonical coordinate projection. Let $\mathcal{F}$ be a coherent sheaf of $\mathcal{O}_X$ -modules whose stalk at the origin in X equals F , and let $\mathcal{G}$ be a coherent $\mathcal{O}_{X^{\{n\}}}$ -module whose stalk at the origin $0^{\{n\}}$ in $X^{\{n\}}$ equals $F^{\tilde{\otimes}_R^n}$. We can identify R with $\mathcal{O}_{Y,0}$ and A with $\mathcal{O}_{X,0}$. Then F is R -flat if and only if $\mathcal{F}_0$ is $\mathcal{O}_{Y,0}$ -flat.

We first prove the “only if” direction of Theorem 1.10, by contradiction. Assume that F is R -flat. Since flatness is an open condition, by Douady’s theorem [5], we can assume that $\mathcal{F}$ and $\mathcal{G}$ are $\mathcal{O}_Y$ -flat. Suppose that $F^{\tilde{\otimes}_R^n}$ has a vertical element over $\mathcal{O}_{Y,0}$. In other words, (after shrinking X and Y if necessary) there exist a nonzero section $\tilde{m} \in \mathcal{G}$ and an analytic subset $Z \subset X^{\{n\}}$, such that $Z_0 = \mathcal{V}(\text{Ann}_{\mathcal{O}_{X^{\{n\}},0^{\{n\}}}}(\tilde{m}_0))$ and the image $\varphi^{\{n\}}(Z)$ has empty interior in Y . Let $\tilde{\varphi}$ denote the restriction $\varphi^{\{n\}}|_Z : Z \rightarrow Y$. Consider $\xi \in Z$ such that the fibre dimension of $\tilde{\varphi}$ at ξ is minimal. Then the fibre dimension $\text{fbd}_x \tilde{\varphi}$ is constant on some open neighbourhood U of ξ in Z . By the Remmert Rank Theorem, $\tilde{\varphi}(U)$ is locally analytic in Y near $\eta = \tilde{\varphi}(\xi)$. Since $\tilde{\varphi}(Z)$ has empty interior in Y , it follows that there is a holomorphic function g in a neighbourhood of η in Y , such that $(\tilde{\varphi}(U))_\eta \subset \mathcal{V}(g_\eta)$. Therefore, $\tilde{\varphi}_\xi^*(g_\eta) \cdot \tilde{m}_\xi = 0$ in $\mathcal{G}_\xi$; i.e., $\mathcal{G}_\xi$ has a (nonzero) zero-divisor in $\mathcal{O}_{Y,\eta}$, contradicting flatness.

We will now prove the more difficult “if” direction of the theorem, by induction on m . If $m = 0$, then F is finitely generated over R , and the result follows from Auslander’s theorem 1.1 (because flatness of finitely generated modules over a local ring is equivalent to freeness, the analytic tensor product equals the ordinary tensor product for finite modules, and vertical elements in finite modules are just zero-divisors).

The inductive step will be divided into three cases:

- (1) F is torsion-free over A ;
- (2) F is a torsion A -module;
- (3) F is neither A -torsion-free nor a torsion A -module.

We will use the following notation for φ and its fibred powers. If $\xi \in X$ and $\eta = \varphi(\xi) \in Y$, then we write $\xi = (\zeta, \eta)$, where $\zeta \in \mathbb{C}^m$. If $\xi_1, \dots, \xi_n \in X$ are all mapped to η , then we denote the point $(\xi_1, \dots, \xi_n) \in X^{\{n\}}$ by $(\bar{\zeta}, \eta)$, where $\bar{\zeta} = (\zeta_1, \dots, \zeta_n) \in \mathbb{C}^{nm}$, $\xi_i = (\zeta_i, \eta)$.

Case (1). We prove this case independently of the inductive hypothesis. We essentially repeat the argument of Galligo and Kwieciński [8], which itself is an adaptation of Auslander [3] to the almost finitely generated context.

Suppose that $F^{\tilde{\otimes}_R^n}$ has no vertical elements over R . Then it follows from Lemma 3.3(1) that $F^{\tilde{\otimes}_R^i}$ has no vertical elements, for $i = 1, \dots, n$.

By Lemma 3.3(3),

$$\mathrm{fd}_R(F^{\tilde{\otimes}_R^n}) = \mathrm{fd}_R(F) + \mathrm{fd}_R(F^{\tilde{\otimes}_R^{n-1}}) = \cdots = n \cdot \mathrm{fd}_R(F).$$

On the other hand, since $F^{\tilde{\otimes}_R^n}$ has no vertical elements over R , it has no zero-divisors over R , so that $\mathrm{depth}_R(F^{\tilde{\otimes}_R^n}) \geq 1$. It follows from Lemma 3.1(2) that $\mathrm{fd}_R(F^{\tilde{\otimes}_R^n}) < n$. Hence $n \cdot \mathrm{fd}_R(F) < n$. This is possible only if $\mathrm{fd}_R(F) = 0$, i.e., F is R -flat.

Case (2). Suppose that F is not R -flat. We will show that then $F^{\tilde{\otimes}_R^n}$ contains vertical elements over R . Let $I = \mathrm{Ann}_A(F)$. Since every element of F is killed by some nonzero element of A , and F is finitely generated over A , then I is a nonzero ideal in A . Put $B = A/I$; then F is finitely generated over B . Let $I(0)$ denote the *evaluation* of I at $y = 0$ (i.e., the result of tensoring I with $\tilde{\otimes}_R R/\mathfrak{m}_R$; $I(0)$ is the ideal generated by I in $A(0) := A\tilde{\otimes}_R R/\mathfrak{m}_R \cong \mathbb{C}\{x_1, \dots, x_m\}$).

First suppose that $I(0) \neq (0)$. Then there exists $g \in I$ such that $g(0) := g(0, x) \neq 0$, and F is a finite $A/(g)A$ -module. It follows that (after an appropriate linear change in the x -coordinates) g is regular in x_m and hence, by the Weierstrass Preparation Theorem, that F is finite over $R\{x_1, \dots, x_{m-1}\}$. Therefore, $F^{\tilde{\otimes}_R^n}$ has a vertical element over R , by the inductive hypothesis.

On the other hand, suppose that $I(0) = (0)$. Then $B(0) \cong \mathbb{C}\{x_1, \dots, x_m\}$. Let Z be a closed analytic subspace of X such that $\mathcal{O}_{Z,0} \cong B$, and let $\tilde{\varphi} := \varphi|_Z$. It follows that the fibre $\tilde{\varphi}^{-1}(0)$ equals $\mathbb{C}^m$. Of course, m is not the generic fibre dimension of $\tilde{\varphi}$, because otherwise all its fibres would equal $\mathbb{C}^m$, so we would have $B = A$ and $I = (0)$, contrary to the choice of I . Therefore, by Proposition 4.2, there is an isolated algebraic vertical component in the n -fold fibred power of $\tilde{\varphi}_0$; i.e., $B^{\tilde{\otimes}_R^n}$ has a zero-divisor in R . But $F^{\tilde{\otimes}_R^n}$ is a finitely generated $B^{\tilde{\otimes}_R^n}$ -module, so itself it has a zero-divisor (hence a vertical element) over R .

Case (3). Suppose that F is not R -flat, F has zero-divisors in A , but $\mathrm{Ann}_A(F) = (0)$. Let

$$K := \{f \in F : af = 0 \text{ for some nonzero } a \in A\};$$

i.e., K is the A -torsion submodule of F . Since K is a submodule of a finitely generated module over a Noetherian ring, K is finitely generated; say $K = \sum_{i=1}^s A \cdot f_i$. Take $a_i \in A \setminus \{0\}$ such that $a_i f_i = 0$, and put $g = a_1 \dots a_s$. Then the sequence of A -modules

$$(5.1) \quad 0 \rightarrow K \rightarrow F \xrightarrow{g} gF \rightarrow 0$$

is exact, and gF is a torsion-free A -module.

First suppose that gF is R -flat. By applying $\tilde{\otimes}_R K$ and $F\tilde{\otimes}_R$ to (5.1), we get exact sequences

$$\begin{aligned} \dots \rightarrow \widetilde{\mathrm{Tor}}_1^R(gF, K) &\rightarrow K\tilde{\otimes}_R K \rightarrow F\tilde{\otimes}_R K \rightarrow gF\tilde{\otimes}_R K \rightarrow 0, \\ \dots \rightarrow \widetilde{\mathrm{Tor}}_1^R(F, gF) &\rightarrow F\tilde{\otimes}_R K \rightarrow F\tilde{\otimes}_R F \rightarrow F\tilde{\otimes}_R gF \rightarrow 0. \end{aligned}$$

But $\widetilde{\mathrm{Tor}}_1^R(gF, K) = 0 = \widetilde{\mathrm{Tor}}_1^R(F, gF)$, since gF is R -flat. So we have injections

$$K\tilde{\otimes}_R K \hookrightarrow F\tilde{\otimes}_R K \hookrightarrow F\tilde{\otimes}_R F,$$

and by induction, an injection $K^{\tilde{\otimes}_R^i} \hookrightarrow F^{\tilde{\otimes}_R^i}$, for all $i \geq 1$. In particular, $K^{\tilde{\otimes}_R^n}$ is a submodule of $F^{\tilde{\otimes}_R^n}$. Since gF is R -flat and F is not R -flat, it follows that K is not R -flat. Therefore, by Case (2), $K^{\tilde{\otimes}_R^n}$ (and hence $F^{\tilde{\otimes}_R^n}$) has a vertical element over R .

Now suppose that gF is not R -flat. We will consider two subcases, depending on whether $g(0) = 0$, where $g(0)$ is the evaluation of g at $y = 0$.

Subcase (3.1). Suppose that $g(0) = 0$. Consider the algebra $B = A/(g)A$. As in Case (2), let Z be a closed analytic subspace of X such that $\mathcal{O}_{Z,0} \cong B$, and let $\tilde{\varphi} = \varphi|_Z : Z \rightarrow Y$. Since $g(0) = 0$ and $B \neq A$, it follows as in Case (2) that the n -fold fibred power of $\tilde{\varphi}_0$ has an isolated algebraic vertical component over Y_0 , by Proposition 4.2. Let (the germ) Z' denote the union of all such components, and let Z'' denote the union of the remaining irreducible components of $Z_{0\{n\}}^{\{n\}}$. Then there exists a nonzero $r \in \mathcal{O}_{Y,0}$ such that Z' is mapped by $\varphi^{\{n\}}$ into the zero-set germ $\mathcal{V}(r)$; hence $rb = 0$ in $B^{\tilde{\otimes}_R^n}$, for every $b \in A^{\tilde{\otimes}_R^n} \cong \mathcal{O}_{X^{\{n\},0\{n\}}}$ which vanishes identically on Z'' when restricted to $Z' \cup Z''$.

Consider the surjection $\rho : F^{\tilde{\otimes}_R^n} \rightarrow (gF)^{\tilde{\otimes}_R^n}$ induced by (5.1). Since gF is a non- R -flat module which is torsion-free over A , then there exists a nonzero vertical element $\bar{m} \in (gF)^{\tilde{\otimes}_R^n}$, by Case (1). Choose $m \in F^{\tilde{\otimes}_R^n}$ such that $\rho(m) = \bar{m}$. Then

$$\mathrm{Ann}_{A^{\tilde{\otimes}_R^n}}(m) \supset \mathrm{Ann}_{A^{\tilde{\otimes}_R^n}}(\bar{m}) \cdot \mathrm{Ann}_{A^{\tilde{\otimes}_R^n}}(B^{\tilde{\otimes}_R^n}),$$

by the definition of ρ . Since $\bar{m}$ is nonzero and vanishes on $Z' \cup Z''$, and $Z'' = \bigcap \mathcal{V}(b)$, where the intersection is over finitely many b as above, we can choose $b \in A^{\tilde{\otimes}_R^n}$ such that both $rb = 0$ in $B^{\tilde{\otimes}_R^n}$ and $b\bar{m} \neq 0$. Then $bm \neq 0$ in $F^{\tilde{\otimes}_R^n}$ (because $\rho(bm) = b\bar{m}$). Furthermore, since $rb = 0$ in $B^{\tilde{\otimes}_R^n}$, we get

$$\mathrm{Ann}_{A^{\tilde{\otimes}_R^n}}(bm) \supset \mathrm{Ann}_{A^{\tilde{\otimes}_R^n}}(\bar{m}) \cdot r.$$

Therefore, $\mathcal{V}(\text{Ann}_{A^{\otimes_R^n}}(bm))$ is mapped into the union of the nowhere-dense germs $\mathcal{V}(r)$ and the image of $\mathcal{V}(\text{Ann}_{A^{\otimes_R^n}}(\bar{m}))$. In other words, bm is a vertical element of $F^{\tilde{\otimes}_R^n}$.

Subcase (3.2). Finally suppose that $g(0) = g(0, x) \neq 0$. We will show that then $(gF)^{\tilde{\otimes}_R^n}$ embeds into $F^{\tilde{\otimes}_R^n}$ (hence so do its vertical elements). The following lemma shows that, in order for $(gF)^{\tilde{\otimes}_R^n}$ to embed into $F^{\tilde{\otimes}_R^n}$, it suffices to prove that $g^{\tilde{\otimes}_R^n}$ is not a zero-divisor of $(gF)^{\tilde{\otimes}_R^n}$.

Lemma 5.1. *Let $R = \mathbb{C}\{y_1, \dots, y_n\}$, and let A and B be regular local analytic R -algebras. Suppose that M and N are finite A - and B -modules (respectively). Let $g \in A$, $h \in B$, and $m \in gM\tilde{\otimes}_R hN$ all be nonzero elements. If $m = 0$ as an element of $M\tilde{\otimes}_R N$, then $g\tilde{\otimes}_R h$ is a zero-divisor of $gM\tilde{\otimes}_R hN$.*

Proof. Using the identification

$$gM\tilde{\otimes}_R hN \cong (gM\tilde{\otimes}_R B) \otimes_{A\tilde{\otimes}_R B} (A\tilde{\otimes}_R hN),$$

we can write $m = \sum_{i=1}^k m_i \otimes n_i$, where the $m_i \in gM\tilde{\otimes}_R B$, and $n_1, \dots, n_k$ generate $A\tilde{\otimes}_R hN$. The latter can be extended to a sequence $n_1, \dots, n_k, n_{k+1}, \dots, n_t$ generating $A\tilde{\otimes}_R N$. Setting $m_{k+1} = \dots = m_t = 0$, we get $m = \sum_{i=1}^t m_i \otimes n_i \in (M\tilde{\otimes}_R B) \otimes_{A\tilde{\otimes}_R B} (A\tilde{\otimes}_R N)$. By [4, Lemma 6.4], $m = 0$ in $M\tilde{\otimes}_R N$ if and only if there are $m'_1, \dots, m'_s \in M\tilde{\otimes}_R B$ and $a_{ij} \in A\tilde{\otimes}_R B$, such that

$$(5.2) \quad \sum_{j=1}^s a_{ij} m'_j = m_i \text{ in } M\tilde{\otimes}_R B, \quad \text{for all } i;$$

$$(5.3) \quad \sum_{i=1}^t a_{ij} n_i = 0 \text{ in } A\tilde{\otimes}_R N, \quad \text{for all } j.$$

Multiplying the equations (5.2) by $g\tilde{\otimes}_R 1$, we get

$$(5.4) \quad \sum_{j=1}^s a_{ij} (g\tilde{\otimes}_R 1) m'_j = (g\tilde{\otimes}_R 1) m_i \text{ in } gM\tilde{\otimes}_R B, \quad \text{for all } i;$$

hence $(g\tilde{\otimes}_R 1)m = 0$ in $gM\tilde{\otimes}_R N$, by (5.3) and (5.4).

Now write $(g\tilde{\otimes}_R 1)m = \sum_{i=1}^l m_i \otimes n_i$, where the $n_i \in A\tilde{\otimes}_R N$, and $m_1, \dots, m_l$ generate $gM\tilde{\otimes}_R B$. Then $(g\tilde{\otimes}_R 1)m = 0$ in $(gM\tilde{\otimes}_R B) \otimes_{A\tilde{\otimes}_R B} (A\tilde{\otimes}_R N)$ if and only if there are $n'_1, \dots, n'_p \in A\tilde{\otimes}_R N$ and $b_{ij} \in A\tilde{\otimes}_R B$,

such that

$$(5.5) \quad \sum_{j=1}^p b_{ij} n'_j = n_i \text{ in } A \tilde{\otimes}_R N, \quad \text{for all } i;$$

$$(5.6) \quad \sum_{i=1}^l b_{ij} m_i = 0 \text{ in } gM \tilde{\otimes}_R B, \quad \text{for all } j.$$

Multiplying the equations (5.5) by $1 \tilde{\otimes}_R h$, we get

$$(5.7) \quad \sum_{j=1}^p b_{ij} (1 \tilde{\otimes}_R h) n'_j = (1 \tilde{\otimes}_R h) n_i \text{ in } A \tilde{\otimes}_R hN, \quad \text{for all } i;$$

hence $(g \tilde{\otimes}_R h)m = 0$ in $gM \tilde{\otimes}_R hN$, by (5.6) and (5.7). Thus $g \tilde{\otimes}_R h$ is a zero-divisor of $gM \tilde{\otimes}_R hN$, as required. $\square$

The next lemma is an immediate consequence of the properties of completed tensor product [16]. Here $\hat{\otimes}_{\mathbb{C}}$ denotes the completed tensor product over $\mathbb{C}$, and $\widehat{\text{Tor}}^{\mathbb{C}}$ denotes its derived functor.

Lemma 5.2. *Let $A \cong \mathbb{C}[[x_1, \dots, x_m]]$, $B \cong \mathbb{C}[[z_1, \dots, z_l]]$, and let $N' \hookrightarrow N$, $M' \hookrightarrow M$ be inclusions of finitely generated A -, B -modules (respectively). Then $N' \hat{\otimes}_{\mathbb{C}} M'$ is a submodule of $N \hat{\otimes}_{\mathbb{C}} M$.*

Proof. Since A and B are faithfully flat over $\mathbb{C}$, $N' \rightarrow N$ and $M' \rightarrow M$ are injective as in the hypotheses if and only if we have monomorphisms of $A \hat{\otimes}_{\mathbb{C}} B$ -modules,

$$0 \rightarrow N' \hat{\otimes}_{\mathbb{C}} B \xrightarrow{\varphi} N \hat{\otimes}_{\mathbb{C}} B \quad \text{and} \quad 0 \rightarrow A \hat{\otimes}_{\mathbb{C}} M' \xrightarrow{\psi} A \hat{\otimes}_{\mathbb{C}} M.$$

The derived functor of the completed tensor product satisfies $\widehat{\text{Tor}}_i^{\mathbb{C}}(\cdot, \cdot) = 0$ for $i \geq 1$ (see [16, p. 104]), so that, by tensoring the first short exact sequence with $A \hat{\otimes}_{\mathbb{C}} M$ and the second with $N' \hat{\otimes}_{\mathbb{C}} B$, we get injections

$$\begin{aligned} \varphi \hat{\otimes}_{\mathbb{C}} 1 &: (N' \hat{\otimes}_{\mathbb{C}} B) \otimes_{A \hat{\otimes}_{\mathbb{C}} B} (A \hat{\otimes}_{\mathbb{C}} M) \hookrightarrow (N \hat{\otimes}_{\mathbb{C}} B) \otimes_{A \hat{\otimes}_{\mathbb{C}} B} (A \hat{\otimes}_{\mathbb{C}} M), \\ 1 \hat{\otimes}_{\mathbb{C}} \psi &: (N' \hat{\otimes}_{\mathbb{C}} B) \otimes_{A \hat{\otimes}_{\mathbb{C}} B} (A \hat{\otimes}_{\mathbb{C}} M') \hookrightarrow (N' \hat{\otimes}_{\mathbb{C}} B) \otimes_{A \hat{\otimes}_{\mathbb{C}} B} (A \hat{\otimes}_{\mathbb{C}} M). \end{aligned}$$

Hence the composite $(\varphi \hat{\otimes}_{\mathbb{C}} 1) \circ (1 \hat{\otimes}_{\mathbb{C}} \psi)$ is injective; i.e., $N' \hat{\otimes}_{\mathbb{C}} M'$ embeds into $N \hat{\otimes}_{\mathbb{C}} M$. $\square$

We can now complete the proof of Subcase (3.2). Since F is a finite A -module, we can write $F = A^q/M$, where $q \geq 1$ and M is a submodule of A^q . Then

$$(g^l F)(0) = g^l(0) \cdot F(0) = g^l(0) \cdot \frac{A^q(0)}{M(0)} = \frac{A^q(0)}{M(0) : g^l(0)},$$

for all $l \geq 1$, where evaluation again means setting $y = 0$ (or, equivalently, taking $\tilde{\otimes}_R R/\mathfrak{m}_R$), and $M(0) : g^l(0) := \{m \in A^q(0) : g^l(0) \cdot m \in M(0)\}$. Since

$$M(0) : g(0) \subset M(0) : g^2(0) \subset \cdots \subset M(0) : g^l(0) \subset \cdots$$

is an increasing sequence of submodules of a Noetherian module $A^q(0)$, it stabilizes; i.e., there exists $k \geq 1$ such that $M(0) : g^{k+1}(0) = M(0) : g^k(0)$. In other words, there exists $k \geq 1$ such that $g(0)$ is not a zero-divisor in $(g^k F)(0)$.

Now let $\mathcal{M}$ be a coherent submodule of $\mathcal{O}_X^q$ such that $\mathcal{F} = \mathcal{O}_X^q/\mathcal{M}$, where $\mathcal{F}_0 = F$, as before, and let $\tilde{g}$ be a section of $\mathcal{O}_X$ in a neighbourhood of the origin, such that the germ of $\tilde{g}$ at 0 equals g . For $l \geq 0$, define

$$\mathcal{M}_l := (\mathcal{M} : \tilde{g}^{l+1})/(\mathcal{M} : \tilde{g}^l).$$

Since $\mathcal{M}_l$ is a finite $\mathcal{O}_X$ -module, it can be written as $\mathcal{M}_l = \mathcal{O}_X^p/\mathcal{N}_l$, where $p \geq 1$ and $\mathcal{N}_l$ is a finite submodule of $\mathcal{O}_X^p$. Then, for every $l \geq 0$, since the condition $(\mathcal{N}_l)_{(\zeta, \eta)}(\eta) = (\mathcal{O}_X^p)_{(\zeta, \eta)}(\eta)$ is open in X , it follows that the vanishing locus of the evaluation of $\mathcal{M}_l$ at points $\eta \in Y$,

$$V_l = \{(\zeta, \eta) \in X : (\mathcal{M}_l)_{(\zeta, \eta)} \tilde{\otimes}_{\mathcal{O}_{Y, \eta}} \mathcal{O}_{Y, \eta}/\mathfrak{m}_{Y, \eta} = 0\}$$

is open in X . Therefore, without loss of generality, we can assume that $\tilde{g}_{(\zeta, \eta)}(\eta)$ is not a zero-divisor of $(\tilde{g}^k \mathcal{F})_{(\zeta, \eta)}(\eta)$, for every (ζ, η) in X .

Finally observe that (because gF is A -torsion-free), multiplication by g induces an isomorphism $gF \rightarrow g^2 F$ of A -modules, and in general, $gF \cong g^l F$, for $l \geq 1$. We thus have isomorphisms $(gF)^{\tilde{\otimes}_R^n} \cong (g^l F)^{\tilde{\otimes}_R^n}$ of $A^{\tilde{\otimes}_R^n}$ -modules, for $l \geq 1$. In particular, for every $l \geq 1$, $g^{\tilde{\otimes}_R^n}$ is a zero-divisor of $(gF)^{\tilde{\otimes}_R^n}$ if and only if it is a zero-divisor of $(g^l F)^{\tilde{\otimes}_R^n}$. To complete the proof of Theorem 1.10 it therefore remains to prove the following:

Lemma 5.3. *Let $k \geq 0$ and suppose that $g(0)$ is not a zero-divisor of $(g^k F)(0)$. Then $g^{\tilde{\otimes}_R^n}$ is not a zero-divisor of $(g^k F)^{\tilde{\otimes}_R^n}$.*

Proof. Suppose the assertion is false. Consider a nonzero $m \in (g^k F)^{\tilde{\otimes}_R^n}$ such that $g^{\tilde{\otimes}_R^n} \cdot m = 0$ in $(g^k F)^{\tilde{\otimes}_R^n}$. Let $\mathcal{G}$ be a coherent $\mathcal{O}_{X^{\{n\}}}$ -module whose stalk at the origin equals $(g^k F)^{\tilde{\otimes}_R^n}$, and let $\tilde{m} \in \mathcal{G}$ be a representative of m . Then the germ $(\tilde{g}_0)^{\tilde{\otimes}_R^n} \cdot \tilde{m}_0$ is zero in $\mathcal{G}_0$; hence

$$\left(\tilde{g}_{(\bar{\zeta}, \eta)}^{\tilde{\otimes}_{\mathcal{O}_{Y, \eta}}^n} \cdot \tilde{m}_{(\bar{\zeta}, \eta)} \right) (\eta) = 0,$$

for all $(\bar{\zeta}, \eta)$ near the origin in $X^{\{n\}}$, where $\bar{\zeta} = (\zeta_1, \dots, \zeta_n)$ and $(\zeta_i, \eta) \in X$, $i = 1, \dots, n$.

On the other hand, we can choose $(\bar{\zeta}, \eta) = (\zeta_1, \dots, \zeta_n, \eta) \in X^{\{n\}}$ arbitrarily close to the origin, so that $\tilde{m}_{(\bar{\zeta}, \eta)}(\eta) \neq 0$. As noted above, we can assume that $\tilde{g}_{(\zeta_i, \eta)}(\eta)$ is not a zero-divisor of $(\tilde{g}^k \mathcal{F})_{(\zeta_i, \eta)}(\eta)$, for $i = 1, \dots, n$. Then, by Lemma 5.2 and faithful flatness of completion, $\tilde{g}_{(\bar{\zeta}, \eta)}^{\tilde{\mathcal{O}}_{Y, \eta}^n}(\eta) = \tilde{g}_{(\zeta_1, \eta)}(\eta) \tilde{\otimes}_{\mathbb{C}} \dots \tilde{\otimes}_{\mathbb{C}} \tilde{g}_{(\zeta_n, \eta)}(\eta)$ is not a zero-divisor of $(\tilde{g}^k \mathcal{F})_{(\bar{\zeta}, \eta)}^{\tilde{\mathcal{O}}_{Y, \eta}^n}(\eta) = (\tilde{g}^k \mathcal{F})_{(\zeta_1, \eta)}(\eta) \tilde{\otimes}_{\mathbb{C}} \dots \tilde{\otimes}_{\mathbb{C}} (\tilde{g}^k \mathcal{F})_{(\zeta_n, \eta)}(\eta)$. Hence

$$\left(\tilde{g}_{(\bar{\zeta}, \eta)}^{\tilde{\mathcal{O}}_{Y, \eta}^n} \cdot \tilde{m}_{(\bar{\zeta}, \eta)} \right) (\eta) \neq 0;$$

a contradiction. $\square$

REFERENCES

1. J. Adamus, *Vertical components in fibre powers of analytic spaces*, J. Algebra **272** (2004), 394–403.
2. J. Adamus, *Flatness testing and torsion freeness of analytic tensor powers*, J. Algebra **289** (2005), 148–160.
3. M. Auslander, *Modules over unramified regular local rings*, Illinois J. Math. **5** (1961), 631–647.
4. D. Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Grad. Texts in Math., Vol. 150, Springer, New York, 1995.
5. A. Douady, *Le problème des modules pour les sous-espaces analytiques compacts d'un espace analytique donné*, Ann. Inst. Fourier (Grenoble) **16:1** (1966), 1–95.
6. G. Fischer, *Complex analytic geometry*, Lecture Notes in Math. **538**, Springer, Berlin, Heidelberg, New York, 1976.
7. J. Frisch, *Points de platitude d'un morphisme d'espaces analytiques complexes*, Invent. Math. **4** (1967), 118–138.
8. A. Galligo and M. Kwieciński, *Flatness and fibred powers over smooth varieties*, J. Algebra **232** (2000), 48–63.
9. H. Grauert and R. Remmert, *Analytische Stellenalgebren*, Springer, Berlin, Heidelberg, New York, 1971.
10. A. Grothendieck, *Eléments de géométrie algébrique IV. Etudes locale des schémas et des morphismes de schémas*, Publ. Math. I.H.E.S. **28**, 1966.
11. H. Hironaka, *Stratification and flatness*, Real and Complex Singularities, Proc. Oslo 1976, ed. Per Holm, Sijthoff and Noordhoff (1977), 199–265.
12. E. Kunz, *Introduction to commutative algebra and algebraic geometry*, Birkhäuser, Basel, Boston, Berlin, 1985.
13. M. Kwieciński, *Flatness and fibred powers*, Manuscripta Mathematica **97** (1998), 163–173.
14. S. Lichtenbaum, *On the vanishing of Tor in regular local rings*, Illinois J. Math. **10** (1966), 220–226.
15. S. Łojasiewicz, *Introduction to Complex Analytic Geometry*, Birkhäuser, Basel, Boston, Berlin, 1991.
16. J.-P. Serre, *Local algebra*, Springer Monographs in Math., Springer, Berlin, 2000.

17. W.V. Vasconcelos, *Flatness testing and torsionfree morphisms*, J. Pure Appl. Algebra **122** (1997), 313–321.
18. W.V. Vasconcelos, Computational methods in commutative algebra and algebraic geometry, Algorithms and Computation in Mathematics, Vol. 2, Springer, Berlin, Heidelberg, 1998.

J. ADAMUS, DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF WESTERN ONTARIO, LONDON, ONTARIO N6A 5B7 CANADA, AND INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, KRAKOW, POLAND

E-mail address: `jadamus@uwo.ca`

E. BIERSTONE, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, M5S 3G3 CANADA

E-mail address: `bierston@math.toronto.edu`

P.D. MILMAN, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, M5S 3G3 CANADA

E-mail address: `milman@math.toronto.edu`